

Jean Mendoza

Dear WA Ecology,
Please see the attached comments from Friends of Toppenish Creek.
Thank you.
Jean Mendoza

Sept. 30, 2022

Dear WA State Dept. of Ecology,

Thank you for including proximity to agriculture as a risk factor for health impacts on overburdened communities.

Remember that there is a wide range of impacts from different types of agriculture. It would be a mistake to paint all forms of agriculture with the same brush. Within Yakima County, according to research conducted by the Friends of Toppenish Creek, air ammonia levels were 63 times higher at a Lower Yakima Valley (LYV) site near concentrated animal feeding operations (CAFOs) compared to an Upper Yakima Valley (UYV) site about sixty miles distant from CAFOs.

Remember that emissions from some aspects of CAFO operations are more serious than others, as we will describe in future comments.

Below is a 2015 listing of research that documents the adverse health impacts from CAFOs.

Sincerely,

Jean Mendoza

Executive Director, Friends of Toppenish Creek

Research Related to Concentrated Animal Feeding Operations and Asthma

American Thoracic Society (1996) Health effects of outdoor air pollution. Committee of the environmental and occupational health assembly of the American Thoracic Society. *Am J Respir Crit Care Med.* 1996 Jan; 153(1):3-50. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8542133>

Aneja VP, Blunden J, Roelle PA, Schlesinger WH, Knighton R, Niyogi D, Gilliam W, Jennings G, Duke CS: Workshop on agricultural air quality: State of the science. *Atmos Environ* 2008, 42 :3195-3208. Retrieved from <http://www.ncsu.edu/airworkshop/>

Chase, L.E. (2011) How much gas do cows produce? *Progressive Dairyman*. Retrieved from http://www.progressivedairy.com/index.php?option=com_content&id=6491:how-much-gas-do-cows-produce&Itemid=121

Barret, J.R. (2006) Hogging the air: CAFO emissions reach into schools. *Environmental Health Perspectives* 114 (4). Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1440820/pdf/ehp0114-a0241a.pdf>

Delfino RJ, Quintana PJ, Floro J, Gastanaga VM, Samimi BS, Kleinman MT, Liu LJS, Bufalino C, Wu CF, McLaren CE: Association of FEV1 in asthmatic children with personal microenvironmental exposure to airborne particulate matter. *Environ Health Perspect* 2007, 115 :1665-1669. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1242025/pdf/ehp0112-000932.pdf>

Dominici F, Peng RD, Zeger SL, White RH, Samet JM: Particulate air pollution and mortality in the United States: did the risks change from 1987 to 2000? *Am J Epidemiol* 2007, 166 :880-888. Retrieved from <http://aje.oxfordjournals.org/content/166/8/880.full.pdf+html>

Kirkhorn S, Garry VF: Agricultural lung diseases. *Environ Health Perspect* 2000, 108 :705-712. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1637683/pdf/envhper00313-0104.pdf>

Koenig JQ, Mar TF, Allen RW, Jansen K, Lumley T, Sullivan JH, Trenga CA, Larson TV, Liu LJS: Pulmonary effects of indoor-and outdoor-generated particles in children with asthma. *Environ Health Perspect* 2005, 113 :499-503. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1278493/>

Koran, H.S. (1995) Associations between criteria air pollutants and asthma. *Environ Health Perspect.* 1995 Sep; 103 Suppl 6:235-42. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1518942/pdf/envhper00365-0226.pdf>

Marchant, J.A., Naleway, A.L., Svendsen, E.T., Kelly, K. M., Burmeister, L.F., Stromquist, A.M., Taylor, C.D., Thorne, P.S., Reynolds, S.J., Sanderson, W. T. & Chrischilles, E.A. (2005) Asthma and farm exposures in a cohort of rural Iowa children. *Environmental Health Perspectives* 113(3): 350 – 356. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1253764/pdf/ehp0113-000350.pdf>

Mirabelli MC, Wing S, Marshall SW, Wilcosky TC. 2006. Asthma symptoms among adolescents who attend public schools that are located near confined swine feeding operations. *Pediatrics*. Jul; 118 (1): e6675. Retrieved from <http://pediatrics.aappublications.org/content/118/1/e66.full.pdf+html>

Omland O: Exposure and respiratory health in farming in temperate zones - A review of the literature. *Ann Agric Environ Med* 2002,9 :119-136. Retrieved from <http://www.aaem.pl/pdf/aaem0219.pdf>

Radon K, Schulze A, Ehrenstein V, van Strien RT, Praml G, Nowak D. 2006. Environmental exposure to confined animal feeding operations and respiratory health of neighboring residents. 18:300-307. Retrieved from http://journals.lww.com/epidem/Fulltext/2007/05000/Environmental_Exposure_to_Confined_Animal_Feeding.4.aspx

Schwartz DA: Etiology and pathogenesis of airway disease in children and adults from rural communities. *Environ Health Perspect* 1999, 107(Suppl 3) :393-401. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1566226/pdf/envhper00520-0013.pdf>

Sigurdarson ST, Kline JN. 2006. School proximity to concentrated animal feeding operations and prevalence of asthma in students. *Chest*. Jun; 129 (6): 1486 – 91. Retrieved from <http://journal.publications.chestnet.org/article.aspx?articleid=1084481>

Von Essen S, Auvermann B: Health effects from breathing air near CAFOs for feeder cattle or hogs. *Journal of Agromedicine* 2005, 10 :55-64. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16702123>

Williams et al (2011) Airborne cow allergen, ammonia and particulate matter at homes vary with distance to industrial scale dairy operations: an exposure assessment. *Environmental Health* 10 :72. Retrieved from <http://www.ehjournal.net/content/10/1/72>

Wilson SM, Serre ML: Examination of atmospheric homes, and schools ammonia levels near hog CAMS, in Eastern North Carolina. *Atmos Environ* 2007, 41 :4977-4987. Retrieved from <http://www.sciencedirect.com/science/article/pii/S1352231007000453>

Zanobetti, A. & Schwartz. J. (2009) The effect of fine and coarse particulate air pollution on mortality: A national analysis. *Environ Health Perspect.* 117(6): 898–903. doi: [10.1289/ehp.0800108](https://doi.org/10.1289/ehp.0800108) Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2702403/pdf/ehp-117-898.pdf>

Research Related to Air in Concentrated Animal Feeding Operations and Public Health

Blanes-Vidal, V., Nadimi, E.S., Ellerman, T., Andersen, H.V., & Lofstrom, P. (2012) Perceived annoyance from environmental odors and association with atmospheric ammonia levels in non-urban residential communities: a cross-sectional study. *Environmental Health* 11(27). Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3458882/pdf/1476-069X-11-27.pdf>

Bowman, A., Mueller, K., & Smith, M. (2000). *Increased animal waste production from concentrated animal feeding operations (CAFOs): Potential implications for public and environmental health*. Nebraska Center for Rural Health Research. Retrieved from <http://www.unmc.edu/rural/documents/cafo-report.pdf>

Bunton B, Shaughnessy P, Fitzsimmons S, Gering J, Hoff S, Lyngbye M, Thorne PS, Wasson J, Werner M: Monitoring and modeling of emissions from concentrated animal feeding operations: Overview of methods. *Environ Health Perspect* 2007, 115:303-307. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1817704/>

Burns, R., Xin, H., Gates, R., Li, H., Hoff, S., Moody, L., et al. (2007). *Tyson broiler ammonia emission monitoring project: Final report*. Retrieved from <http://www.sierraclub.org/environmentallaw/lawsuits/docs/ky-tysonreport.pdf>

Center for Livable Future. (2009). *Flies may spread drug-resistant bacteria from poultry operations*. Retrieved from <http://www.jhsph.edu/news/news-releases/2009/graham-flies.html>

Cole D, Todd L, Wing S: Concentrated swine feeding operations and public health: A review of occupational and community health effects. *Environ Health Perspect* 2000, 10:685-699. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1638284/pdf/envhper00309-0041.pdf>

Donham KJ, Wing S, Osterberg D, Flora JL, Hodne C, Thu KM, Thorne PS: Community health and socioeconomic issues surrounding concentrated feeding operations. *Environ Health Perspect* 2007, 115 :317-320. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1817697/pdf/ehp0115-000317.pdf>

Donham, K.J. (2002) Synergistic effects of dust and ammonia on the occupational health effects of poultry production workers. *Journal of Agromedicine* 8 (2). Retrieved from <http://openagricola.nal.usda.gov/Record/IND23302625>

Donham KJ, Lee JA, Reynolds SJ. (2006). Assessment of air quality at neighbor residences in the vicinity of swine production facilities. *Journal of Agromedicine*. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19274894>

Government Accountability Office. (2008). *Concentrated animal feeding operations: EPA needs more information and a clearly defined strategy to protect air and water quality from pollutants of concern*. Retrieved from <http://www.gao.gov/new.items/d08944.pdf>

Gilchrist, M.J., Greko, C., Wallinga, D.B., Beran, G.W., Riley, D.G. & Thorne, P.S. (2007) The potential role of concentrated animal feeding operations in infectious disease epidemics and antibiotic resistance. *Environmental Health Perspectives* 115 (2): 313-316. doi: 10.1289/ehp.8837. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1817683/>

Homes MJ, Heber AJ, Wu LK: Viability of bioaerosols produced from a swine facility. Retrieved from <http://www.extension.purdue.edu/pork/house/conf.htm>

Godbout, S., Lemay, S.P., Duchaine, C., Pelletier, F., Larouche, J.P., Belzile, M., & Feddes, J.J. (2009) Swine production impact on residential ambient air quality. *Journal of Agromedicine* 14(3): 291 – 298. doi: 10.1080/10599230903041943. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19657878>

Green CG, Gibbs SG, Tarwater PM, Mota LC, Scarpino PV: Bacterial plume emanating from the air surrounding swine confinement operations. *J Occ Environ Hygiene* 2006, 3 :9-15. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16482973>

Heederik D, Sigsgaard T, Thorne PS, Kline JN, Avery R, Bonlokke JH, Chrischilles EA, Dosman JA, Duchaine C, Kirkhorn SR, Kulhankova K, Merchant JA: Health effects of airborne exposures from concentrated animal feeding operations. *Environ Health Perspect* 2007, 115 :298-302. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1817709/pdf/ehp0115-000298.pdf>

Horton, R.A., Wing, S., Marshall, S.W., & Brownley, K.A. (2009) Malodor as a trigger of stress and negative mood in neighbors of industrial hog operations. *Am J Public Health*. 2009 November; 99(S3): S610–S615. doi: [10.2105/AJPH.2008.148924](https://doi.org/10.2105/AJPH.2008.148924). Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2774199/pdf/S610.pdf>

Kirkhorn, S.R. (2002) Community and environmental health effects of concentrated animal feeding operations. *Minn. Med. Oct.*: 85(10): 38-43. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12416314>

Kleinman, M. (2000). *The health effects of air pollution on children*. Retrieved from http://www.aqmd.gov/forstudents/health_effects_on_children.pdf

Kullman GJ, Thorne PS, Waldron PF, Marx JJ, Ault B, Lewis DM, Siegel PD, Olenchock SA, Merchant JA: Organic dust exposures from work in dairy barns. *Am Ind Hyg Assoc J* 1998, 59 :403-413. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9670470>

Michigan Department of Environmental Quality (MDEQ) Toxics Steering Group (TSG). (2006). *Concentrated animal feedlot operations (CAFOs) chemicals associated with air emissions*. Retrieved from http://www.michigan.gov/documents/CAFOs_Chemicals_Associated_with_Air_Emissions_5-10-06_158862_7.pdf

McConnell, P. E., Coggins, M. A., Hogan, V.A., & Flemming, G.T. (2008) Exposure assessment of airborne contaminants in the indoor environment of Irish swine farms. *Annals of Agricultural Environmental Medicine* 15(2): 323 – 6. Retrieved from <http://www.aaem.pl/pdf/15323.pdf>

Mirabelli, M.C., Wing, S., Marshall, S.W., & Wilcosky, T.C. (2006). Race, poverty, and potential exposure of middle-school students to air emissions from confined swine feeding operations. *Environmental Health Perspectives*, 114(4), 591–596. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1440786/pdf/ehp0114-000591.pdf>

[National Association of Local Boards of Public Health \(2010\) Understanding Confined Animal Feeding Operations and Their Impact on Communities](http://www.cdc.gov/nceh/ehs/docs/understanding_cafos_nalboh.pdf). Retrieved from http://www.cdc.gov/nceh/ehs/docs/understanding_cafos_nalboh.pdf

Oliver DM, Clegg CD, Heathwaite AL, Haygarth PM: Preferential attachment of *Escherichia coli* to different particle size fractions of an agricultural grassland soil. *Water Air Soil Pollution* 2007, 185 :369-375. Retrieved from http://eprints.lancs.ac.uk/26432/1/Corrected_Oliver_et_al_Ecoli_attach_wasp.pdf

Pew Commission on Industrial Animal Farm Production. (2009). *Putting meat on the table: Industrial farm animal production in America*. Retrieved from <http://www.ncifap.org/images/PCIFAPFin.pdf>

Predicala BZ, Maghirang RG: Field comparison of inhalable and total dust samplers for assessing airborne dust in swine confinement barns. *Appl Occup Environ Hyg* 2003, 18 :694-701. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12909537>

Purdy CW, Clark RN, Straus DC: Analysis of aerosolized particulates of feedyards located in the southern high plains of Texas. *Aerosol Science and Technology* 2007, 41 :497-509. Retrieved from <http://www.cprl.ars.usda.gov/wmru/pdfs/Analysis%20of%20aerosolized%20particulates%20of%20feed%20yards%20located%20in%20the%20Southern%20High%20Plains%20of%20TX.pdf>

Purdy CW, Clark RN, Straus DC: Ambient and indoor particulate aerosols generated by dairies in the southern high plains. *J Dairy Sci* 2009, 92 :6033-6045. Retrieved from <http://download.journals.elsevierhealth.com/pdfs/journals/0022-0302/PIIS0022030209713207.pdf>

Reynolds SJ, Donham KJ, Stookes-berry J. 1997. Air quality assessments in the vicinity of swine production facilities. *Journal of Agromedicine*. 4:37-46. Retrieved from http://www.tandfonline.com/doi/abs/10.1300/J096v04n01_06#preview

Schenker MB, Christiani D, Cormier Y, Dimich-Ward H, Doekes G, Dosman JA: American Thoracic Society: respiratory health hazards in agriculture. *Am J Resp Crit Care Med* 1998, 158 :S1-S76. Retrieved from <http://www.thoracic.org/statements/resources/eoh/agriculture1-79.pdf>

Schiffman SS, Studwell CE, Landerman LR, Berman K, Sundy JS. 2005. Symptomatic effects of exposure to diluted air samples from a swine confinement atmosphere on health human subjects. *Environmental Health Perspectives*. 113:567-76. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1257549/pdf/ehp0113-000567.pdf>

Schmalzried, H.D. & Fallon, L.F., Jr. (2007). Large-scale dairy operations: Assessing concerns of neighbors about quality-of-life issues. *Journal of Dairy Science*, 90(4), 2047-2051. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17369247>

Thorne P: Environmental health impacts of concentrated animal feeding operations: anticipating hazards – searching for solutions. *Environ Health Perspect* 2007, 115 :296-297. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1817701/pdf/ehp0115-000296.pdf>

Thorne, P. (2007) Environmental health impacts of concentrated animal feeding operations: Anticipating hazards – searching for solutions. *Environmental Health Perspectives*. 115(2). Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1817701/pdf/ehp0115-000296.pdf>

Thu K, Donham K, Ziegenhorn R, Reynolds S, Thorne PS, Subramanian P, Whitten P, Stookesberry J: A control study of the physical and mental health of residents living near a large-scale swine operation. *J Agric Saf Health* 1997, 3 :13-26. Retrieved from <http://www.sraproject.org/wp-content/uploads/2007/12/acontrolstudyofthephysicalandmentalhealth.pdf>

Thu KM: Public health concerns for neighbors of large-scale swine production operations. *J Agric Saf Health* 2002, 8 :175-184. Retrieved from <http://www.dnr.missouri.gov/env/apcp/docs/phc-swine-thu.pdf>

Wall, S.K. & Ebner, P. (2007) CAFOs and public health: emissions and the respiratory health of neighbors. *Purdue University Extension Service*. Retrieved from <http://www.extension.purdue.edu/extmedia/ID/cafo/ID-358-W.pdf>

Wing, S. & Wolf, S. (2000) Intensive livestock operations, health, and quality of life among eastern North Carolina residents, *Environ Health Perspect*. 2000 March; 108(3): 233–238. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1637983/pdf/envhper00304-0089.pdf>

Wing, S., Horton, R.A. & Rose, K.M. (2013) Air pollution from industrial swine operations and blood pressure of neighboring residents. *Environmental Health Perspectives* 2013 Jan;121(1):92-6. doi: 10.1289/ehp.1205109. Epub 2012 Oct 18. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3553433/>

Wing S, Horton RA, Marshall SW, Thu K, Tajik M, Schinasi L, Schiffman SS: Air pollution and odor in communities near industrial swine operations. *Environ Health Perspect* 2008, 116: 1362-1368. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2569096/>

Research related to Asthma, Allergy & Cow Dander:

Gupta S, Bidani RK, Jhamb S, Agarwal MK. Role of animal danders as inhalant allergens in bronchial asthma in India. *J Asthma* 1996;33(5):339-48 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8827941>

Hinze S, Bergmann K, Lowenstein H, Hansen G: Cow hair allergen (Bos d 2) content in house dust: Correlation with sensitization in farmers with cow hair asthma. *Int Arch Allergy Imm* 1997, 112 :231-237. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9066508>

Holopainen E, Salo OP, Tarkianen E, Malmberg H. The most important allergens in allergic rhinitis. *Acta Otolaryngol* 1979;Suppl 360;16-18 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/287333>

Kanerva L, Estlander T. Immediate and delayed skin allergy from cow dander. *Am J Contact Dermat* 1997;8(3):167-9 Retrieved from http://journals.lww.com/dermatitis/Abstract/1997/09000/Immediate_and_Delayed_Skin_Allergy_From_Cow_Dander.9.aspx

Kanerva L, Susitaival P. Cow dander: the most common cause of occupational contact urticaria in Finland. *Contact Dermatitis* 1996 Nov;35(5):309-10 Retrieved from <http://onlinelibrary.wiley.com/doi/10.1111/j.1600-0536.1996.tb02400.x/abstract>

Kanerva L, Toikkanen J, Jolanki R, Estlander T. Statistical data on occupational contact urticaria. *Contact Dermatitis* 1996 Oct;35(4):229-33 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8957643>

Mantylarvi R, Parkkinen S, Rytkonen M, Pentikainen J, Pelkonen J, Rautiainen J, Zeiler T, Virtanen T. Complementary DNA cloning of the predominant allergen of bovine dander: a new member in the lipocalin family. *J Allergy Clin Immunol* 1996;97(6):1297-303 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8648026>

Prahl P. Allergens in cow hair and dander. Origin of cow allergens in the environment. *Allergy* 1981;36(8):561-71 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7337202>

Prahl P, Sonderstrup Hansen G. Lymphocyte responses to an aqueous extract from cow hair and dander. Investigations of asthmatics and normal controls with lymphocyte transformation tests. *Allergy* 1982;37(3):155-60 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/6182813>

Prahl P, Bucher D, Plesner T, Weeke B, Lowenstein H. Isolation and partial characterisation of three major allergens in an extract from cow hair and dander. *Int Arch Allergy Appl Immunol* 1982;67(4):293-301 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7068248>

Rautiainen J, Pentikainen J, Rytkonen M, Linnala-Kankkunen A, Pelkonen J, Virtanen T, Mantylarvi R. Molecular analysis of allergenic proteins in bovine dander. *Allergy* 1996;51(6):378-82 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8837659>

Rautiainen J, Ryttonen M, Syrjanen K, Pentikainen J, Zeiler T, Virtanen T, Mantyjarvi R. Tissue localization of bovine dander allergen Bos d 2. *J Allergy Clin Immunol* 1998;101(3):349-53 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9525451>

Rautiainen J, Auriola S, Konttinen A, Virtanen T, Ryttonen-Nissinen M, Zeiler T, Mantyjarvi R. Two new variants of the lipocalin allergen Bos d 2. *J Chromatogr B Biomed Sci Appl* 2001;763(1-2):91-8 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11710588>

Rautalahti M, Terho EO, Vohlonen I, Husman K. Atopic sensitization of dairy farmers to work-related and common allergens. *Eur J Respir Dis Suppl* 1987;152:155-64 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/3478214>

Santa H, Saarela JT, Laatikainen R, Rautiainen J, Virtanen T, Ryttonen M, Mantyjarvi R. A bovine dander allergen, comparative modeling, and similarities and differences in folding with related proteins. *J Protein Chem* 1998;17(7):657-62 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9853680>

Spiewak R. Sensitization to cow and pig allergens among farmers in Eastern Poland. [Polish] *Med Pr* 2001;52(5):351-4 Retrieved from <http://www.radoslawspiewak.net/2001-5a.htm>

Szepfalusi Z, Ebner C, Urbanek R, Ebner H, Scheiner O, Boltz-Nitulescu G, Kraft D. Detection of IgE antibodies specific for allergens in cow milk and cow dander. *Int Arch Allergy Immunol* 1993;102(3):288-94 Retrieved from <http://www.karger.com/Article/Abstract/236538>

Terho EO, Husman K, Vohlonen I, Rautalahti M, Tukiainen H. Allergy to storage mites or cow dander as a cause of rhinitis among Finnish dairy farmers. *Allergy* 1985;40(1):23-6 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/3977027>

Valero Santiago AL, Rosell Vives E, Lluch Perez M, Sancho Gomez J, Piulats Xanco J, Malet Casajuana A. Occupational allergy caused by cow dander: detection and identification of the allergenic fractions. *Allergol Immunopathol (Madr)* 1997;25(6):259-65 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9469201>

Virtanen, T., Zeiler, T., Rautiainen, J., Taivanainen, A., Pentikainen, J., Ryttonen, M., Parkkinen, S., Pelkonen, J., Mantyjarvi. R. (2006) Immune reactivity of cow-asthmatic dairy farmers to the major allergen of cow (BDA20) and to other cow-derived proteins. The use of purified BDA20 increases the performance of diagnostic tests in respiratory cow allergy. *Clinical & Experimental Allergy* 26 (2). pp. 188 – 196. DOI: 10.1111/j.1365-2222.1996.tb00079.x Retrieved from <http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2222.1996.tb00079.x/abstract>

Ylonen J, Mantyjarvi R, Taivainen A, Virtanen T. Comparison of the antigenic and allergenic properties of three types of bovine epithelial material. *Int Arch Allergy Immunol* 1992;99(1):112-7 Retrieved from <http://www.karger.com/Article/Pdf/236343>

Ylonen J, Mantyjarvi R, Taivainen A, Virtanen T. IgG and IgE antibody responses to cow dander and urine in farmers with cow-induced asthma. *Clin Exp Allergy* 1992;22(1):83-90 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/1551038>

Zeiler T, Taivainen A, Mantyjarvi R, Tukiainen H, Rautiainen J, Rytkonen-Nissinen M, Virtanen T: Threshold levels of purified natural Bos d 2 for inducing bronchial airway response in asthmatic patients. *Journal of the British Society for Allergy and Clinical Immunology* 2002, 32: 1454-1460. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12372125>

Research Related to Air Quality and Animal Health

Chang, C.W., Chung, H., Huang, C.F. & Su, H.J. (2001) Exposure assessment to airborne endotoxin, dust, ammonia, hydrogen sulfide and carbon dioxide in open style swine houses. *Annals of Occupational Hygiene* 45 (6): 457 – 65. Retrieved from <http://annhyg.oxfordjournals.org/content/45/6/457.full.pdf>

Curtis S E, Anderson CR, Simon J, Jensen AH, Day DL, Kelley KW. 1975. Effects of aerial ammonia, hydrogen sulfide and swine – house dust on rate of gain and respiratory tract structure in swine. *J Anim Sci* 41(3):735 Retrieved from <http://www.journalofanimalscience.org/content/41/3/735.full.pdf>

Done, S.H., Chennells, D.J., Gresham, A.C., Williamson, S., Taylor, L.L., Bland, V., Jones, P., Armstrong, D., White, R.P., Demmers, T.G., Teer, N. & Wathes, C.M. (2005) Clinical and pathological responses of weaned pigs to atmospheric ammonia and dust. *Veterinary Research* 157 (3): 71 – 80. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16024672>

Donham KJ. 2000. The Concentration of Swine Production. *Veterinary Clinics of North America: Food Animal Practice*. 16:559-597. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11084993>

Donham, K.J. (1991) Association of environmental air contaminants with disease and productivity in swine. *American Journal of Veterinary Research* 52 (10). Retrieved from <http://openagricola.nal.usda.gov/Record/IND92016191>

Donham, K.J. (1989) Relationships of air quality and productivity in intensive swine housing. *Agri-Practice* 10 (6). Retrieved from <http://openagricola.nal.usda.gov/Record/IND90031609>

Drummond JG, Curtis SE, Simon J. 1978. Effects of atmospheric ammonia on pulmonary bacterial clearance in the young pig. *Am J Vet Res*. 39:211 –212. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/343668>

Drummond JG, Curtis SE, Simon J, Norton HW. 1980. Effects of aerial ammonia on growth and health of young pigs. *J Anim Sci* 50(6):1085 –1091. Retrieved from <http://www.journalofanimalscience.org/content/50/6/1085.full.pdf>

Drummond JG, Curtis SE, Meyer RC, Simon J, Norton HW. 1981a. Effects of atmospheric ammonia on young pigs experimentally infected with *Bordetella bronchiseptica*. *Am J Vet Res* 42(6):963 –968. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7283246>

Drummond JG, Curtis SE, Simon J, Norton HW. 1981b. Effects of atmospheric ammonia on young pigs experimentally infected with *Ascaris suum*. *Am J Vet Res* 42(6): 969 – 974. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7283247>

Environmental Health Sciences Research Center (2002) Iowa Concentrated Animal Feeding Operations Air Quality Study. Retrieved from <http://www.public-health.uiowa.edu/ehsrc/cafostudy.htm>

Gurk S, Brunsch R, Kaufman O. 1997. Systematic analysis of trace gases in dairy cow housing. *Landtechnik* 52: 206 - 207.

Hamilton TDC, Roe JM, Webster AF. 1996. The synergistic role of gaseous ammonia in the aetiology of *Pasterella multocida* induced atrophic rhinitis in swine. *J Clin Microbiol* 43:2185 – 2190.

Hamilton, T.D., Roe, J.M., Hayes, C.M., Jones, P., Pearson, G.R., & Webster, A.J. (1999) Contributory and exacerbating roles of gaseous ammonia and organic dust in the etiology of atrophic rhinitis. *Clin Diagn Lab Immunol* 6(2): 199 – 203. Retrieved from <http://cvi.asm.org/content/6/2/199.long>

Harper. L.A., Sharpe, R.R., & Simmons, J.D. (2004) Ammonia emissions from swine houses in the southeastern United States. *Journal of Environmental Quality* 33(2): 449 – 57. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15074795>

Hauser RH, Folsch DW. 1988. Methods for measuring atmospheric ammonia in poultry houses: review and approved practices. *Journal of Veterinary Medicine* 35: 579 – 593. Retrieved from <http://japr.fass.org/content/14/3/622.full.pdf>

Heber, A.J., Lim, T.T., Ni, J.Q., Tao, P.C., Schmidt, A.M., Koziel, J.A., Hoff, S.J., Jacobson, L. D., Ahzng, Y., Baughman, G.B. (2006) Quality assured measurements of animal building emissions: particulate matter concentrations. *Journal of Animal Waste Management* 56 (12): 1842 – 8. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17195483>

Hoff, S.J., Bundy, D.S., Nelson, M.A., Zelle, B. C., Jacobson, L.D., Heber, A.J., Ni, J., Zhang, Y., Koziel, J.A., & Beasley, D.B. (2006) Emissions of ammonia, hydrogen sulfide, and odor before, during, and after slurry removal from a deep-pit swine finisher. *Journal of Air Waste Management Assoc.* 56(5): 581 – 90. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16739794>

[Holland, R.E., Carson, T.L. & Drummond, K.J. Animal Health Effects. *Environmental Health Sciences Research Center. Iowa Study of CAFO s* Retrieved from \[http://www.ehsrc.uiowa.edu/CAFOstudy/CAFO_6-2.pdf\]\(http://www.ehsrc.uiowa.edu/CAFOstudy/CAFO_6-2.pdf\)](http://www.ehsrc.uiowa.edu/CAFOstudy/CAFO_6-2.pdf)

Kim, K.Y., Ko, H.J., Lee, K.J., Park, J.B., & Kim, C.N. (2005) Temporal and spatial distributions of aerial contaminants in an enclosed pig building in winter. *Environmental Research* 99(2): 150 – 7. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16194664>

Kim, K.Y., Ko, H.J., Kim, H.T., Kim, Y.S., & Roh, Y.M. (2008) Effect of manual feeding on the level of farmer's exposure to airborne contaminants in the confinement nursery pig house. *Industrial Health (NIOSH)* 46 (2): 138 – 43. Retrieved from https://www.jstage.jst.go.jp/article/indhealth/46/2/46_2_138/pdf

Klenz RD, Fedde MR. 1978. Hydrogen sulfide: effects on avian respiratory control and intrapulmonary carbon dioxide receptors. *Respir Physiol* 32(3):355 – 358. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/208128>

Malayer, J.R., Kelly, D.T., Diekman, M.A., Brandt, K.E., Sutton, A.L., Long, G.G., & Jones, D.D. (1987) Influence of manure gases on puberty in gilts. *Journal of Animal Science* 64(5): 1476 – 83. Retrieved from <http://www.journalofanimalscience.org/content/64/5/1476.full.pdf>

O'Shaughnessy, P.T., Achutan, C. & Karsten, A.W. (2002) Temporal variation of indoor air quality in an enclosed swine confinement building. *Journal of Agricultural Safety and Health*. 8(4): 349 – 64. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12549241>

Pickrell, J. (1991) Hazards in confinement housing – gases and dusts in confined animal houses for swine, poultry, horses and humans. *Veterinary Human Toxicology* 33 (1): 32 – 9. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/2017863>